

Computer Graphics

Ágoston Sipos
siposagoston@inf.elte.hu

Eötvös Loránd University
Faculty of Informatics

2025-2026. Fall semester

Introduction and contact details

- ▶ Lecturer: Ágoston Sipos
- ▶ E-mail: `siposagoston@inf.elte.hu`
- ▶ Room: D 2.610
- ▶ Consultation: after a discussion by email
 - ▶ Preferred time: Monday between 12 and 15
- ▶ Information on the Canvas

Table of Contents

Motivation

Coordinate-systems

- Cartesian coordinate system

- Polar coordinate system

- Barycentric coordinates

- Homogeneous coordinates


Modelling

- ▶ How do we represent our virtual world?
 - ▶ How do we represent the points of our virtual space, how do we store it on the computer?
 - Coordinate systems
 - ▶ How do we represent the simple geometric building blocks (line, plane, triangle, etc.)?
 - Set of points
 - Description in different coordinate systems


Points, vectors

- ▶ Point: an element of the Euclidean plane/space that has no extent whatsoever.
- ▶ Vector:
 - ▶ algebraically: element of a vector space.
 - ▶ geometrically: a displacement that has *direction* and *magnitude*
 - ▶ additional operations interpreted on vectors: addition, subtraction, multiplication by a scalar, cross product (result is a vector), dot product (result is a scalar)

Addition



Subtraction



Dot product

Let two vectors be, $\mathbf{a} = [a_x, a_y, a_z]$ and $\mathbf{b} = [b_x, b_y, b_z]$. Dot product is denoted by $\langle \mathbf{a}, \mathbf{b} \rangle$, i.e


$$\langle \mathbf{a}, \mathbf{b} \rangle = a_x b_x + a_y b_y + a_z b_z.$$

This can also be expressed as

$$\langle \mathbf{a}, \mathbf{b} \rangle = |\mathbf{a}| \cdot |\mathbf{b}| \cdot \cos(\alpha),$$

where α is the angle between $\mathbf{a}$ and $\mathbf{b}$ vectors.

Cross product (3D)



- ▶ $\mathbf{a} \times \mathbf{b}$ is perpendicular to both $\mathbf{a}$ and $\mathbf{b}$
- ▶ $\mathbf{a}$, $\mathbf{b}$ and $\mathbf{a} \times \mathbf{b}$: right-hand rule



Cross product of vectors

Cross product as a determinant:

$$\begin{aligned}\begin{bmatrix} a_x \\ a_y \\ a_z \end{bmatrix} \times \begin{bmatrix} b_x \\ b_y \\ b_z \end{bmatrix} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix} \\ &= \mathbf{i} \cdot \begin{vmatrix} a_y & a_z \\ b_y & b_z \end{vmatrix} - \mathbf{j} \cdot \begin{vmatrix} a_x & a_z \\ b_x & b_z \end{vmatrix} + \mathbf{k} \cdot \begin{vmatrix} a_x & a_y \\ b_x & b_y \end{vmatrix} \\ &= \begin{bmatrix} a_y b_z - a_z b_y \\ -a_x b_z + a_z b_x \\ a_x b_y - a_y b_x \end{bmatrix}\end{aligned}$$

Points, vectors

- ▶ Point and vector can be represented with coordinates of a chosen coordinate system. BUT: let's pay attention to the operations that can be performed!
- ▶ Operations between points and vectors:
 - ▶ point + vector = point displaced point
 - ▶ point - point = vector difference vector
 - ▶ point + point not interpreted!



Notation

- ▶ Points: $\mathbf{a} \in \mathbb{E}^2, \mathbf{b} \in \mathbb{E}^3, \dots$
- ▶ Vectors: $\mathbf{v} \in \mathbb{R}^n, n = 2, 3, \dots$
 - ▶ special: $[\mathbf{v}]_0 \in \mathbb{R}^n$ is a vector that is unit long i.e.
 $||[\mathbf{v}]_0||_2 = 1$.
- ▶ Lines: $e, f, g, \dots$
- ▶ Planes: $S, \dots$
- ▶ Matrices: $\mathbf{M} \in \mathbb{R}^{n \times m}$

Coordinate-system

- ▶ Uniquely represents point of space with n-tuples

$$\text{pl.: } \mathbf{p} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \in \mathbb{E}^3$$

- ▶ It allows storing points on the computer
- ▶ It allows the use of algebraic and analytical tools to solve geometric problems
- ▶ It may be easier to describe a problem in a coordinate system that fits it well


Cartesian coordinate system

- ▶ Descartes, 1637.: *dissertation on the method* (Discours de la méthode pour bien conduire sa raison et chercher la vérité dans les sciences)
- ▶ Most of the time we come across this, this is the easiest and most common way of representing points

Cartesian coordinate system


- ▶ For each finite point of the Euclidean space [plane] we assign an ordered, real number triplet (x, y, z) [pair (x, y)]
- ▶ We define Cartesian coordinate system with a starting point (origin, $\mathbf{o}$) in the space, and an orthonormal system: three pairwise perpendicular unit vectors: $\mathbf{i}, \mathbf{j}$ and $\mathbf{k}$ (these specify the direction of the x, y, z axes).
- ▶ Then the coordinates x, y, z of a point $\mathbf{p}$ in row are the same as the signed orthogonal projections of the vector $\mathbf{p} - \mathbf{o}$ to the orthonormal basis vectors $\mathbf{i}, \mathbf{j}, \mathbf{k}$.
- ▶ *Reminder:* the signed orthogonal projection of the vector $\mathbf{a}$ on unit vector $[\mathbf{b}]_0$ is $\langle \mathbf{a}, [\mathbf{b}]_0 \rangle = |\mathbf{a}| \cos \angle(\mathbf{a}, [\mathbf{b}]_0)$

Signed orthogonal projection



Geometric interpretation

- *More conceptually:* $\mathbf{p}(a, b, c)$ is the point from the origin which we get by moving a units along the x axis, then b units along the y axis, and finally c units along z .





Geometric interpretation

- ▶ According to the interpretation above, using the unit-long base vectors $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ pointing in the direction of the coordinate axes, $[a, b, c]^T$ the coordinates describe the following point:



$$\mathbf{p} = \mathbf{o} + a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$$

$$= \mathbf{o} + a \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + b \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + c \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$



Orientation – right-handed system



Orientation – left-handed system



Orientation – left-handed system





Calculating distance

- ▶ The distance between two points **a** and **b** is d if:
$$d^2 = (b_1 - a_1)^2 + (b_2 - a_2)^2$$
- ▶ This is due to Pitagoras's theorem: the sum of the squared sides of the right-angled triangle are equal
- ▶ $\rightarrow$ this opens up the possibility of describing new shapes (e.g. circle, sphere)
- ▶ Generally:
$$\mathbf{a}, \mathbf{b} \in \mathbb{E}^n : d = \sqrt{\langle \mathbf{b} - \mathbf{a}, \mathbf{b} - \mathbf{a} \rangle} = \sqrt{\sum_{i=1}^n (b_i - a_i)^2}$$
- ▶ Length of a vector: $\mathbf{v} \in \mathbb{R}^n : \|\mathbf{v}\|_2 = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle} = \sqrt{\sum_{i=1}^n v_i^2}$

Planar polar coordinate system

- ▶ Is defined by a starting point **o** (reference point) and a half-line starting from it (polar axis).
- ▶ The location of a point **p** is determined by two data: (r, θ)
 - ▶ $r \geq 0$: is the distance between **p** and **o**
 - ▶ $\theta \in [0, 2\pi)$: is the angle between the polar axis and the half-line starting from **o** and going towards **p**



Conversions

► Polar $\rightarrow$ Cartesian: $(r, \theta) \rightarrow (x, y)$

► $x = r \cos \theta$

► $y = r \sin \theta$

► Cartesian $\rightarrow$ Polar: $(x, y) \rightarrow (r, \theta)$

► $r = \sqrt{x^2 + y^2}$



$$\theta = \begin{cases} \operatorname{arctg}\left(\frac{y}{x}\right), & x > 0 \wedge y \geq 0 \\ \operatorname{arctg}\left(\frac{y}{x}\right) + 2\pi, & x > 0 \wedge y < 0 \\ \operatorname{arctg}\left(\frac{y}{x}\right) + \pi, & x < 0 \\ \frac{\pi}{2}, & x = 0 \wedge y > 0 \\ \frac{3\pi}{2}, & x = 0 \wedge y < 0 \end{cases}$$
$$= \operatorname{atan2}(y, x)$$

Conversions

- ▶ The above is true if the Cartesian origin and the polar reference point, respectively the Cartesian x-axis and the polar axis, are the same.
- ▶ But what if $x = 0, y = 0$? In this case, with $r = 0$, we get back the origin with an arbitrary angle! The polar angle is not clear, we check whether $r = 0$ before trying to use the conversion formulas from above


Remark

- ▶ We usually use it when it fits well with the things we want to depict, e.g. circular motion
- ▶ Drawbacks: moving from one polar coordinate system (PCS) to another is expensive, calculating derivatives is expensive, ...


Spherical coordinates

- ▶ Spherical coordinate system; defined by a base plane (and its PCS) and a perpendicular "Z axis"
- ▶ A point $\mathbf{p}$ in space is represented by three data: (r, θ, ϕ)
 - ▶ (ϱ, ϕ) : polar coordinates of the projection of $\mathbf{p}$ onto the base plane
 - ▶ $\theta \in [0, \pi]$: the angle between Z axis and the half-line from $\mathbf{o}$ towards $\mathbf{p}$
 - ▶ r : the distance between $\mathbf{p}$ and origin (if $r = 0$ then once again the two polar angles can be anything! This must be checked before conversions)

Spherical coordinates



Spherical coordinates



Conversions

- ▶ Under conditions similar to the planar:
- ▶ Spherical $\rightarrow$ Cartesian: $(r, \theta, \phi) \rightarrow (x, y, z)$

$$x = r \sin \theta \cos \phi,$$

$$y = r \sin \theta \sin \phi,$$

$$z = r \cos \theta$$

- ▶ Cartesian $\rightarrow$ Spherical: $(x, y, z) \rightarrow (r, \theta, \phi)$

$$r = \sqrt{x^2 + y^2 + z^2}$$

$$\phi = \text{atan2}(y, x), \quad r \neq 0$$

$$\theta = \arccos \frac{z}{r}, \quad r \neq 0$$

Remark

- ▶ It is useful, for example, for identifying points on the earth's surface (but there $\theta \in [-\pi/2, \pi/2]$).
- ▶ The parametric representation of a sphere or ellipsoid also utilizes spherical coordinates

Barycentric coordinates


- ▶ August Ferdinand Möbius [1827]
- ▶ Motivation: often only a specific, finite part of the space is interesting to us. We are looking for a representation more "balanced" than the Cartesian representation.

Motivation: intervals



What u, v weights should we place at the ends of the rod if we want the rod to stay in balance when elevated at the point denoted by a triangle?

Motivation: intervals



- ▶ It does not tilt if $(x - a)v = (b - x)u$, where x is the position of the triangle.
- ▶ Only the ratio of u, v is bound by the above, let us further assume that $u + v = 1$
- ▶ Then the weights should be:

$$u = \frac{x - a}{b - a}, v = \frac{b - x}{b - a}$$

Center of mass

- ▶ Mechanical analogy: center of mass for a point system
- ▶ Let us have 3 points in the plane and place weight $m_i \in \mathbb{R}$ in each $\mathbf{p}_i$ point. Then the center of mass is:

$$\mathbf{m} = \sum_{i=0}^2 \frac{m_i}{\sum_{i=0}^n m_i} \mathbf{p}_i$$

- ▶ *Homogeneous* representation: multiplying the weights by a number $h \neq 0$ gives the same point.

Barycentric coordinates

- If $\mathbf{a}_0, \dots, \mathbf{a}_n$ points in $\mathbb{E}^n$ span the space (that is, they do not fall into an $n - 1$ dimensional subspace), then for any $\mathbf{x}$ point of the space we can find $\lambda_0, \dots, \lambda_n$ real numbers that uniquely represents it such that

$$\mathbf{x} = \sum_{i=0}^n \lambda_i \mathbf{a}_i,$$


where λ_i barycentric coordinates satisfy that,

$$\sum_{i=0}^n \lambda_i = 1.$$

Remark

- ▶ In a plane, you need 3 affinely independent points (those that do not fall either in a straight line or a point), in space 4 affinely independent points
- ▶ If $\forall i \lambda_i \geq 0$, then we are talking about a convex combination and the result falls inside the convex hull of the points
- ▶ Affine transformations don't change barycentric coordinates (see later)

Planar barycentric coordinate system



Barycentric $\rightarrow$ Cartesian conversion

- ▶ Let (u, v, w) be the barycentric coordinates of a point and $\mathbf{p}_1 = (x_1, y_1)$, $\mathbf{p}_2 = (x_2, y_2)$, $\mathbf{p}_3 = (x_3, y_3) \in \mathbb{E}^2$ affinely independent points.
- ▶ Then the Cartesian coordinates of the point $\mathbf{x}(x, y)$ represented by (u, v, w) are $\mathbf{x} = u\mathbf{p}_1 + v\mathbf{p}_2 + w\mathbf{p}_3$, i.e.

$$x = ux_1 + vx_2 + wx_3$$

$$y = uy_1 + vy_2 + wy_3$$

Planar barycentric coordinate system

- ▶ Let $\Delta(\mathbf{a}, \mathbf{b}, \mathbf{c}) := \begin{vmatrix} 1 & 1 & 1 \\ a_x & b_x & c_x \\ a_y & b_y & c_y \end{vmatrix}$, $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{E}^2$
- ▶ $\Delta(\mathbf{a}, \mathbf{b}, \mathbf{c})$ equals to twice the signed area of the triangle bound by $\mathbf{a}, \mathbf{b}, \mathbf{c}$ points (positive if the vertices are given in counter-clockwise direction, otherwise it's negative)
- ▶ If we are in $\mathbb{E}^3$: $\Delta(\mathbf{a}, \mathbf{b}, \mathbf{c}) = \langle (\mathbf{b} - \mathbf{a}) \times (\mathbf{c} - \mathbf{a}), \mathbf{n} \rangle$, where $\mathbf{n}$ is the unit long normal of the 3 point's plane.

Cartesian $\rightarrow$ barycentric conversion

- Let $\mathbf{x} \in \mathbb{E}^2$ be a point, then its barycentric coordinates with $\mathbf{p}_1 = (x_1, y_1)$, $\mathbf{p}_2 = (x_2, y_2)$, $\mathbf{p}_3 = (x_3, y_3) \in \mathbb{E}^2$ affinely independent points are:


$$u = \frac{\Delta(\mathbf{x}, \mathbf{p}_2, \mathbf{p}_3)}{\Delta(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3)}$$

$$v = \frac{\Delta(\mathbf{p}_1, \mathbf{x}, \mathbf{p}_3)}{\Delta(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3)}$$

$$w = \frac{\Delta(\mathbf{p}_1, \mathbf{p}_2, \mathbf{x})}{\Delta(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3)}$$

Motivation

- Project the points of a line e onto the x axis from k point!



Motivation

- ▶ The point $\mathbf{d}'$ is not on the Euclidean plane ($\mathbb{E}^2$), since the projection ray passing through $\mathbf{k}$ and $\mathbf{d}$ is parallel to the x axis
- ▶ Idea: expand $\mathbb{E}^2$!
- ▶ $\rightarrow$ Let us consider the same orientation of the lines (their direction) as a point!
- ▶ This will be the *ideal point* of the line.

Definition – ideal point

- ▶ Line = Line + 1 ideal point such that:
 - ▶ Parallel lines have the same ideal point ("they meet at infinity")
 - ▶ The ideal points of a plane lie on a line, this is the *ideal line of the plane*
 - ▶ The ideal line of parallel planes coincide
 - ▶ The ideal elements of space (points, lines) lie in a plane, this is the *ideal plane of the space*

Definition and properties – homogeneous space

- ▶ Projective plane: the projective closure of $\mathbb{E}^2$, that is all the points of $\mathbb{E}^2$ and its ideal line
 - ▶ Two points determine a line in the projective plane
 - ▶ Two lines determine a point (!)
 - ▶ ...
- ▶ Projective space: the projective closure of $\mathbb{E}^3$, that is $\mathbb{E}^3$ plus its ideal plane
 - ▶ Three points determine a plane
 - ▶ Three distinct planes determine a point (!)
 - ▶ (HW: is it true that *any* (arbitrary) three planes define a point in projective space that is on all three planes? In what cases does it not?)
 - ▶ ...

Homogeneous coordinates

- ▶ We assign a quadrilateral *homogeneous coordinates* for each point of the Euclidean space:


$$\begin{aligned}\mathbf{p}(x, y, z) &\rightarrow [x, y, z, 1] \\ &\approx h[x, y, z, 1] \\ &= [hx, hy, hz, h], \quad h \neq 0\end{aligned}$$

- ▶ for every direction vector $\mathbf{v} = [x, y, z]^T$:

$$\begin{aligned}[x, y, z] &\rightarrow [x, y, z, 0] \\ &\approx h[x, y, z, 0] \\ &= [hx, hy, hz, 0], \quad h \neq 0\end{aligned}$$

- ▶ Thus the homogeneous coordinates of a point or direction vector can be multiplied with any $h \neq 0$ number and the resulting coordinates represent the same point or direction vector.
 - ▶ When embedding $\mathbb{E}^n$ into $\mathbb{R}^{n+1}$, this means that all points on the projecting line mean the same euclidean point/vector.

Embedding $\mathbb{E}^2$ into $\mathbb{R}^3$



Converting back to Cartesian coordinate system

- ▶ What does $[x_1, x_2, x_3, x_4]$ represent in projective space?
 - ▶ If $x_4 \neq 0$, then we are talking about a point whose coordinates after homogeneous (or projective) division are:

$$[x_1, x_2, x_3, x_4] \approx \left[\frac{x_1}{x_4}, \frac{x_2}{x_4}, \frac{x_3}{x_4}, 1 \right] = \mathbf{p} \left(\frac{x_1}{x_4}, \frac{x_2}{x_4}, \frac{x_3}{x_4} \right)$$

- ▶ If $x_4 = 0$, but $x_1^2 + x_2^2 + x_3^2 \neq 0$ (=not all zero), then its an ideal point of a line whose orientation is same as $[x_1, x_2, x_3]$ vector.
 - ▶ If $x_i = 0$, $i = 1, 2, 3, 4$, then its undefined.

Notable homogeneous points

- ▶ Let $c \neq 0$ real number. Examples of notable points:
 - ▶ $[0, 0, 0, c]$ origin
 - ▶ $[c, 0, 0, 0]$ ideal point of x axis
 - ▶ $[0, c, 0, 0]$ ideal point of y axis
 - ▶ $[0, 0, c, 0]$ ideal point of z axis

Properties

- ▶ On the projective plane, point and line, and in projective space, point and plane are dual concepts
- ▶ Note that some properties do not transfer from Euclidean space:
 - ▶ A point on a line does not divide the line into two parts! But: two different points do
 - ▶ A line does not divide a plane into two parts! But: two different lines do
 - ▶ Two points do not uniquely identify a segment! (The ideal point of the line "glues together" the two ends of the line)